

Optical tug-of-war tweezers: shaping light for dynamic control of bacterial cells-corrigendum

Joshua Lamstein¹, Anna Bezryadina^{1,2}, Daryl Preece³, Joseph C. Chen⁴,
and Zhigang Chen^{1,5,*}

¹*Department of Physics and Astronomy, San Francisco State University, San Francisco, CA 94132, USA*

²*Department of Electrical and Computer Engineering, University of California San Diego, La Jolla, CA 92093, USA*

³*Department of NanoEngineering, University of California San Diego, La Jolla, CA 92093, USA*

⁴*Department of Biology, San Francisco State University, San Francisco, CA 94132, USA*

⁵*The MOE Key Laboratory of Weak-Light Nonlinear Photonics, and TEDA Applied Physical Institute and School of Physics, Nankai University, Tianjin 300457, China*

**Corresponding author: zhigang@sfsu.com*

The authors would like to apologize for an error in the paper Chinese Optics Letters vol. 15, no. 3, page 030010.

On page 030010-3, the caption for Figs. 5 (a)–(c) should read “2 μm silica” (not 3 μm polystyrene).

doi: 10.3788/COL201715.093502.

References

1. J. Lamstein, A. Bezryadina, D. Preece, J. C. Chen, and Z. Chen, Chin. Opt. Lett. **15**, 030010 (2017).